



# Analysis of Molecular Dynamics Simulations of Biomolecules

- A very complicated arrangement of hundreds of groups interacting with each other
- Where to start to look at?
- What to analyze?
- How much can we learn from simulations?


It is very important to get acquainted with your system

# Aquaporins

## Membrane water channels



# Lipid Bilayer Permeability



**Water** is an exception:

- Small size
- Lack of charge
- Its high concentration

# Water Transport Across Cell Membrane

Always passive; bidirectional; osmosis-driven

- Diffusion through lipid bilayers

slower, but enough for many purposes


- Channel-mediated

Large volumes of water needed to be transported  
(kidneys).

Fast adjustment of water concentration is necessary  
(RBC, brain, lung).


# The Aquaporin Superfamily

Discovered  
in 1992




# Aquaporins in Human Body

Aquaporin-0	Eye: lens fiber cells	Fluid balance of the lens
Aquaporin-1	Red blood cells Kidney: proximal tubules Eye: ciliary epithelium Brain: choroid plexus Lung: alveolar	Osmotic protection Concentration of urine Aqueous humor Production of CSF Alveolar hydration
Aquaporin-2	epithelial cells Kidney: collecting ducts	ADH hormone activity
Aquaporin-3	Kidney: collecting ducts Trachea: epithelial cells	Reabsorption of water Secretion of water
Aquaporin-4	Kidney: collecting ducts Brain: ependymal cells Brain: hypothalamus Lung: bronchial	Reabsorption of water CSF fluid balance Osmosensing function?
Aquaporin-5	epithelium Salivary glands Lacrimal glands	Bronchial fluid Production of saliva secretion Production of tears
Aquaporin-6	Kidney	Very low water permeability!
Aquaporin-7	Testis and sperm	
Aquaporin-8	Testis, pancreas, liver	
Aquaporin-9	Leukocytes	
Aquaporin-10		





Additional members are suspected to exist.


# Aquaporins in the Kidney



# High Permeation to Water



>200 Liters  
Water  
Everyday!



Aquaporins of known structure:



**GlpF** - E. coli glycerol channel (aquaglycerolporin)

**AQP1** - Mammalian aquaporin-1 (pure water channel)


AqpZ and AQPO (2004)

# Functionally Important Features

- Tetrameric architecture
- Amphipathic channel interior
- Water and glycerol transport
- Protons, and other ions are excluded
- Conserved asparagine-proline-alanine residues; NPA motif
- Characteristic half-membrane spanning structure





# A Semi-hydrophobic channel



# Molecular Dynamics Simulations

Protein: ~ 15,000 atoms  
Lipids (POPE): ~ 40,000 atoms  
Water: ~ 51,000 atoms  
Total: ~ 106,000 atoms



NAMD, CHARMM27, PME

NpT ensemble at 310 K


1ns equilibration, 4ns production

10 days /ns - 32-proc Linux cluster


3.5 days/ns - 128 O2000 CPUs

**0.35 days/ns - 512 LeMieux CPUs**

# Protein Embedding in Membrane




Hydrophobic  
surface of the  
protein




Ring of  
Tyr and Trp

# Embedding GlpF in Membrane




# GlpF in VMD




# A Recipe for Membrane Protein Simulations

- Insert your protein into a hydrated lipid bilayer.
- Fix the protein; minimize the rest and run a short "constant-pressure" MD to bring lipids closer to the protein and fill the gap between the protein and lipids.
- Watch water molecules; if necessary apply constraints to prevent them from penetrating into the open gaps between lipids and the protein.
- Monitor the volume of your simulation box until it is almost constant. Do not run the system for too long during this phase.
- Now release the protein, minimize the whole system, and start an NpT simulation of the whole system.
- If desired, you may switch to an NVT simulation, when the system reaches a stable volume.

# Lipid-Protein Packing During the Initial NpT Simulation






# Adjustment of Membrane Thickness to the Protein Hydrophobic Surface





# An extremely stable protein

## Stability of NPA - NPA Interaction




# Glycerol-Saturated GlpF





# Description of full conduction pathway



# Complete description of the conduction pathway

Constriction region



} Selectivity filter

# Details of Protein-Substrate Interaction Are Always Important


- Identify those groups of the protein that are directly involved in the main function of the protein.
- Look at the interaction of these primary residues with other groups in the protein.
- Look at buried charged residues inside the protein; they must have an important role.
- Backbone hydrogen bonds are mainly responsible for stabilization of secondary structure elements in the protein; side chain hydrogen bonds could be functionally important.

# Channel Hydrogen Bonding Sites

...


```
{set frame 0}{frame < 100}{incr frame}{  
  animate goto $frame  
  set donor [atomselect top  
    "name O N and within 2 of  
    (resname GCL and name HO)"]  
  lappend [$donor get index] list1  
  set acceptor [atomselect top  
    "resname GCL and name O and  
    within 2 of (protein and name HN HO)"]  
  lappend [$acceptor get index] list2  
}
```

...




# Channel Hydrogen Bonding Sites

GLN	41	OE1 NE2	LEU	197	O
TRP	48	O NE1	THR	198	O
GLY	64	O	GLY	199	O
ALA	65	O	PHE	200	O
HIS	66	O ND1	ALA	201	O
LEU	67	O	ASN	203	ND2
ASN	68	ND2			
ASP	130	OD1	LYS	33	HZ1 HZ3
GLY	133	O	GLN	41	HE21
SER	136	O	TRP	48	HE1
TYR	138	O	HIS	66	HD1
PRO	139	O N	<u>ASN</u>	68	HD22
ASN	140	OD1 ND2	TYR	138	HN
HIS	142	ND1	ASN	140	HN HD21 HD22
THR	167	OG1	HIS	142	HD1
GLY	195	O	GLY	199	HN
PRO	196	O	<u>ASN</u>	203	HN HD21HD22
			<u>ARG</u>	206	HE HH21HH22





# Channel Hydrogen Bonding Sites

GLN	41	OE1 NE2	LEU	197	O
TRP	48	O NE1	THR	198	O
GLY	64	O	GLY	199	O
ALA	65	O	PHE	200	O
HIS	66	O ND1	ALA	201	O
LEU	67	O	ASN	203	ND2
ASN	68	ND2			
ASP	130	OD1	LYS	33	HZ1 HZ3
GLY	133	O	GLN	41	HE21
SER	136	O	TRP	48	HE1
TYR	138	O	HIS	66	HD1
PRO	139	O N	<u>ASN</u>	68	HD22
ASN	140	OD1 ND2	TYR	138	HN
HIS	142	ND1	ASN	140	HN HD21 HD22
THR	167	OG1	HIS	142	HD1
GLY	195	O	GLY	199	HN
PRO	196	O	<u>ASN</u>	203	HN HD21HD22
			<u>ARG</u>	206	HE HH21HH22




# The Substrate Pathway is formed by C=O groups



# The Substrate Pathway is formed by $C=O$ groups


Non-helical motifs  
are stabilized by  
two **glutamate**  
residues.




# Conservation of Glutamate Residue in Human Aquaporins

```



AQP0_HUMAN ---LNTLHPAVSVGQATTVEIFLTLOFVLCIFATYDE-RRNGQLG
AQP1_HUMAN ---RNDLADGVNSGQGLGIEIIGTLQLVLCVLATDR-RRRDLGG
AQP2_HUMAN ---VNALSNSTTAGQAVTVELFLTLOLVLCIFASTDE-RRGENPG
AQP3_HUMAN GIFATYPSGHLDMINGFFDQFIGTASLIVCVLAIVDPYNNPVPRG
AQP4_HUMAN ---VTMVHGNLTAGHGLLVELLITFQLVFTIFASCDS-KRTDVTG
AQP5_HUMAN ---VNALNNNTTQGOAMVELLITFQLALCIFASTDS-RRTSPVG
AQP6_HUMAN ---INVVRNSVSTGOAVAVELLELTLOLVLCVFASTDS-RQTS--G
AQP7_HUMAN GIFATYLPDHMTLWRGFINEAWLTGMLQLCLFAITDQENNPALPG
AQP8_HUMAN -AAFVTVQEQGQVAGALVAEILTLLALAVCMGAIN--EKTGKP
AQP9_HUMAN HIFATYPAPYLSLANAFADQVATMILLIIVFAIFDSRNLGAPRG
GLPF_ECOLI  GTFSTYPNPHINFVQAFVEMVITAILMGLLILALTDDGNGVPRGP
ruler      ...180.....190.....200.....210.....220
    
```




# Glycerol - water competition for hydrogen bonding sites




# Revealing the Functional Role of Reentrant Loops




Potassium channel



# Single Glycerol per channel





**initial state**






Note that glycerols moved, but not as extensively as earlier!

**final (1ns) state**





**We need to enforce an entire conduction event.**

# Steered Molecular Dynamics






# SMD Simulation of Glycerol Passage




Trajectory of glycerol pulled by **constant force**

# Evidence for Stereoselectivity of Glycerol



Cannot be verified by experimental measurements

# Free Energy Calculation in SMD



Free energy

SMD simulation  
a **non-equilibrium** process

$$\Delta G \leq \langle W \rangle$$

One needs to discount  
irreversible work

$$e^{-\Delta G / k_B T} = \left\langle e^{-W / k_B T} \right\rangle$$

Jarzynski, *PRL* 1997

Hummer, *PNAS*, *JCP* 2001

Liphardt, et al., *Science* 2002

# Constructing the Potential of Mean Force


4 trajectories

$v = 0.03, 0.015 \text{ \AA/ps}$


$k = 150 \text{ pN/\AA}$

$$f(t) = -k[z(t) - z_0 - vt]$$

$$W(t) = \int_0^t dt' v f(t')$$




# Features of the Potential of Mean Force




- Captures major features of the channel
- The largest barrier  $\approx$  **7.3 kcal/mol**; **exp.:  $9.6 \pm 1.5$  kcal/mol**

# Glycerol-Free GlpF




# Water permeation




18 water conducted  
In 4 monomers in 4 ns  
1.125 water/monomer/ns  
Exp. =  $\sim 1-2$  /ns



5 nanosecond  
Simulation




7-8 water  
molecules in  
each channel



# Correlated Motion of Water in the Channel





**The single file of water molecules is maintained.**





# Correlated Motion of Water in the Channel

The single file of water molecules is maintained.






# Diffusion of Water in the channel



One dimensional diffusion:  $2Dt = \langle (z_t - z_0)^2 \rangle$


Experimental value for AQP1:  $0.4-0.8 \text{ e-}5$

# Diffusion of Water in the channel




Improvement of statistics



## Density of O and H atoms along the GlpF channel





# Water Distribution in Aquaporins




# Water Bipolar Configuration in Aquaporins



# Water Bipolar Configuration in Aquaporins



**channel region (20 Å)**





## R E M E M B E R:


One of the most useful advantages of simulations over experiments is that you can modify the system as you wish: You can do modifications that are not even possible at all in reality!

This is a powerful technique to test hypotheses developed during your simulations. **Use it!**

# Electrostatic Stabilization of Water Bipolar Arrangement




# Proton transfer through water




# Cl<sup>-</sup> channel


Anti-parallel



Parallel (barrel stave)



# K<sup>+</sup> channel



# Aquaporins

# Proton Blocking by a Global Orientation Mechanism

