



# Carbon Nanotubes

## Hydrophobic channels - Perfect Models for Membrane Water Channels



A balance between the size and hydrophobicity


# Water-nanotube interaction can be easily modified



Modifying charges  
Modifying vdW parameters

# Carbon Nanotubes


## Hydrophobic channels - Perfect Models for Membrane Water Channels



- Much better statistics
- No need for membrane and lipid molecules



# Carbon Nanotubes

## Hydrophobic channels - Perfect Models for Membrane Water Channels




- Much better statistics
- No need for membrane and lipid molecules

# Water Single-files in Carbon Nanotubes



**Water files form polarized chains in nanotubes**

# Calculation of Diffusion Permeability from MD




$\Phi_0$ : number of water molecules crossing the channel from the left to the right in unit time

$$P_d = \frac{V_w}{N_A} \Phi_0$$


$\Phi_0$  can be directly obtained through **equilibrium MD** simulation by counting “full permeation events”

Simulation of osmotic pressure induced water transport may be done by adding salt to one side of the membrane.



There is a small problem with this setup!


Problem: The solvents on the two sides of a membrane in a conventional periodic system are connected.







We can include more layers of membrane and water to create two compartment of water that are not in contact

NaCl



# UNIT CELL



Semipermeable membrane

Semipermeable membrane


# Realizing a Pressure Difference in a Periodic System

$$P_1 = P_2 + nf \Rightarrow \Delta P = nf / A$$

Fangqiang Zhu


$f$  is the force on each water molecule, for  $n$  water molecules

The overall translation of the system is prevented by applying constraints or counter forces to the membrane.



# Applying a Pressure Difference Across the Membrane

$$\Delta P = nf / A$$




Applying  
force on all  
water  
molecules.

Not a good  
idea!

# Applying a Pressure Difference Across the Membrane

$$\Delta P = nf / A$$




Applying  
force on  
bulk water  
only.

Very good

# Applying a Pressure Difference Across the Membrane

$$\Delta P = n f / A$$




Applying force only on a slab of water in bulk.

Excellent


$P_f$  can be calculated from these simulations

$$\Phi_w = P_f A \left( \frac{\Delta P}{RT} - \Delta C_s \right)$$

# Calculation of osmotic permeability of water channels



$p_f: 1.4 \times 10^{-13} \text{ cm}^3/\text{s}$



$p_f: 7.0 \pm 0.9 \times 10^{-14} \text{ cm}^3/\text{s}$

Exp:  $5.4 - 11.7 \times 10^{-14} \text{ cm}^3/\text{s}$

# Interactive Molecular Dynamics

**VMD** ←·····→ **NAMD**

