




The Computational Microscope

Computational microscope views at atomic resolution ...




... how living cells maintain health and battle disease

John Stone


Our Microscope is Made of...

Chemistry




$$U(\vec{R}) = \underbrace{\sum_{bonds} k_i^{bond} (r_i - r_0)^2}_{U_{bond}} + \underbrace{\sum_{angles} k_i^{angle} (\theta_i - \theta_0)^2}_{U_{angle}} + \underbrace{\sum_{dihedrals} k_i^{dih} [1 + \cos(n_i \phi_i + \delta_i)]}_{U_{dihedral}} + \underbrace{\sum_i \sum_{j \neq i} 4 \epsilon_{ij} \left[\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^6 \right]}_{U_{nonbond}} + \sum_i \sum_{j \neq i} \frac{q_i q_j}{\epsilon r_{ij}}$$

NAMD Software



Virus




Physics

$$m_i \frac{d^2 \vec{r}_i}{dt^2} = \vec{F}_i = -\vec{\nabla} U(\vec{R})$$

Math

$$\vec{r}_i(t + \Delta t) = 2\vec{r}_i(t) - \vec{r}_i(t - \Delta t) + \frac{\Delta t^2}{m_i} \vec{F}_i(t)$$

(repeat **one billion times** = microsecond)



..and Supercomputers

NAMD impact is broad and deep

- Comprehensive, industrial-quality software
 - Integrated with VMD for simulation setup and analysis
 - Portable extensibility through Tcl scripts (also used in VMD)
 - Consistent user experience from laptop to supercomputer
- Large user base – 51,000 registered users
 - 9,100 (18%) are NIH-funded; many in other countries
 - 14,100 have downloaded more than one version
- Leading-edge simulations
 - “most-used software” on NICS Cray XT5 (largest NSF machine)
 - “by far the most used MD package” at TACC (2nd and 3rd largest)
 - NCSA Blue Waters early science projects and acceptance test
 - Argonne Blue Gene/Q early science project



BTRC for Macromolecular Modeling and Bioinformatics
<http://www.ks.uiuc.edu/>


Beckman Institute, UIUC

Outside researchers choose NAMD and succeed


Corringer, et al., *Nature*, 2011

2100 external citations since 2007

Voth, et al., *PNAS*, 2010




180K-atom 30 ns study of anesthetic binding to bacterial ligand-gated ion channel provided “complementary interpretations...that could not have been deduced from the static structure alone.”



Bound Propofol Anesthetic

500K-atom 500 ns investigation of effect of actin depolymerization factor/cofilin on mechanical properties and conformational dynamics of actin filament.




Bare actin

Cofilactin

Recent NAMD Simulations in *Nature*

- M. Koeksal, et al., *Taxadiene synthase structure and evolution of modular architecture in terpene biosynthesis*. (2011)
- C.-C. Su, et al., *Crystal structure of the CusBA heavy-metal efflux complex of Escherichia coli*. (2011)
- D. Slade, et al., *The structure and catalytic mechanism of a poly(ADP-ribose) glycohydrolase*. (2011)
- F. Rose, et al., *Mechanism of copper(II)-induced misfolding of Parkinson's disease protein*. (2011)
- L. G. Cuello, et al., *Structural basis for the coupling between activation and inactivation gates in K(+) channels*. (2010)
- S. Dang, et al., *Structure of a fucose transporter in an outward-open conformation*. (2010)
- F. Long, et al., *Crystal structures of the CusA efflux pump suggest methionine-mediated metal transport*. (2010)
- R. H. P. Law, et al., *The structural basis for membrane binding and pore formation by lymphocyte perforin*. (2010)
- P. Dalhaimer and T. D. Pollard, *Molecular Dynamics Simulations of Arp2/3 Complex Activation*. (2010)
- J. A. Tainer, et al., *Recognition of the Ring-Opened State of Proliferating Cell Nuclear Antigen by Replication Factor C Promotes Eukaryotic Clamp-Loading*. (2010)
- D. Krepiy, et al., *Structure and hydration of membranes embedded with voltage-sensing domains*. (2009)
- N. Yeung, et al., *Rational design of a structural and functional nitric oxide reductase*. (2009)
- Z. Xia, et al., *Recognition Mechanism of siRNA by Viral p19 Suppressor of RNA Silencing: A Molecular Dynamics Study*. (2009)

The Molecular Dynamics Simulation Process



For textbooks see:

M.P. Allen and D.J. Tildesley. *Computer Simulation of Liquids*. Oxford University Press, New York, 1987.

D. Frenkel and B. Smit. *Understanding Molecular Simulations. From Algorithms to Applications*. Academic Press, San Diego, California, 1996.

A. R. Leach. *Molecular Modelling. Principles and Applications*. Addison Wesley Longman, Essex, England, 1996.

More at <http://www.biomath.nyu.edu/index/course/99/textbooks.html>

Classical Dynamics at 300K

Energy function: $U(\vec{r}_1, \vec{r}_2, \dots \vec{r}_N) = U(\vec{R})$

used to determine the force on each atom:

$$m_i \frac{d^2 \vec{r}_i}{dt^2} = \vec{F}_i = -\vec{\nabla} U(\vec{R})$$

yields a set of 3N coupled 2nd-order differential equations that can be propagated forward (or backward) in time.

Initial coordinates obtained from crystal structure, velocities taken at random from Boltzmann distribution.

Maintain appropriate temperature by adjusting velocities.

Classical Dynamics

discretization in time for computing

$$m_i \frac{d^2 \vec{r}_i}{dt^2} = \vec{F}_i = -\vec{\nabla} U(\vec{R})$$

Use positions and accelerations at time t and the positions from time $t-\delta t$ to calculate new positions at time $t+\delta t$.

$$\begin{aligned} \mathbf{r}(t + \delta t) &\approx \mathbf{r}(t) + \mathbf{v}(t)\delta t + \frac{1}{2}\mathbf{a}(t)\delta t^2 \\ \mathbf{r}(t - \delta t) &\approx \mathbf{r}(t) - \mathbf{v}(t)\delta t + \frac{1}{2}\mathbf{a}(t)\delta t^2 \end{aligned} \quad +$$

“Verlet algorithm”






$$-\vec{\nabla} U(\vec{R}) / m_i$$

$$\mathbf{r}(t + \delta t) \approx 2\mathbf{r}(t) - \mathbf{r}(t - \delta t) + \mathbf{a}(t)\delta t^2$$

Potential Energy Function of Biopolymer

- Simple, fixed algebraic form for every type of interaction.
- Variable parameters depend on types of atoms involved.

$$U(\vec{R}) = \underbrace{\sum_{bonds} k_i^{bond} (r_i - r_0)^2}_{U_{bond}} + \underbrace{\sum_{angles} k_i^{angle} (\theta_i - \theta_0)^2}_{U_{angle}} + \underbrace{\sum_{dihedrals} k_i^{dihe} [1 + \cos(n_i \phi_i + \delta_i)]}_{U_{dihedral}} + \underbrace{\sum_i \sum_{j \neq i} 4\epsilon_{ij} \left[\left(\frac{\sigma_{ij}}{r_{ij}} \right)^{12} - \left(\frac{\sigma_{ij}}{r_{ij}} \right)^6 \right] + \sum_i \sum_{j \neq i} \frac{q_i q_j}{\epsilon r_{ij}}}_{U_{nonbond}}$$



heuristic

from physics

Improving the Force Field

- Atomic polarizability increases computation by 2x...
- ...but, the additional computations are perfectly suited to the GPU!
- For now, NAMD calculates atomic polarizability on CPUs only...soon we will also use GPUs

Atomic polarizability of water, highly accurately simulated through additional particles (shown in green)



Molecular Dynamics Ensembles

Constant energy, constant number of particles (NE)

Constant energy, constant volume (NVE)

Constant temperature, constant volume (NVT)

Constant temperature, constant pressure (NPT)

Choose the ensemble that best fits your system and start the simulations, but use NE to check on accuracy of the simulation.

Langevin Dynamics

for temperature control

Langevin dynamics deals with each atom separately, balancing a small friction term with Gaussian noise to control temperature:

$$m \ddot{\vec{r}} = \vec{F}(\vec{r}) - \gamma m \dot{\vec{r}} + \vec{R}(t)$$

$$\langle \vec{R}(t) \cdot \vec{R}(t') \rangle = 6k_B T \gamma \delta(t - t')$$

Langevin Dynamics

for pressure control

***Underlying Langevin-Hoover barostat equation for all atoms:
Equations solved numerically in NAMD***

$$\frac{d^2 V(t)}{dt^2} = \frac{1}{W_{bs}} [P(t) - P_{\text{target}}] - \frac{1}{\tau_{bs}} \frac{dV(t)}{dt} + R_{bs}(t)$$

$$P = \rho k_B T + \frac{1}{Vd} \sum_{i < j} \langle r_{ij} \frac{dU_{\text{tot}}(r_{ij})}{dr_{ij}} \rangle \quad d = \text{dimension}$$


$$\langle R_{bs}(t) R_{bs}(t') \rangle = \frac{2 k_B T_{\text{target}} \delta(t - t')}{W_{bs} \tau_{bs}} \quad W_{bs} = d N_{\text{atoms}} k_B T_{\text{target}} \tau_{\text{period}}^2$$

$$\dot{\mathbf{r}}_i = \mathbf{v}_i + s \mathbf{r}_i \quad \dot{\mathbf{v}}_i = \mathbf{F}_i / m_i - s \mathbf{v}_i$$

$$\dot{V} = dV_s \quad \dot{s} = dV(P - P_{\text{target}}) / W - s / \tau_{bs} + R(t)$$

d - dimension


All-Atom Molecular Dynamics Extends Today to Small Cellular Organelle (100nm)³ Scale



From 10,000 to 100,000 Atom MD in 2000


100k atom MD reached in 2000

- then a factor 10 increase in computation;
- needed to describe membrane processes;
- was achieved through cluster computing;
- produced good quality results for aquaporin;
- is now standard.



E. Tajkhorshid,
P. Nollert,
M. Jensen,
L. Miercke,
J. O'Connell,
and K. Schulten.
Science, **296**:525-530,
2002.


100,000 atoms, 12 ns




F. Khalili-Araghi, V. Jogini,
V. Yarov-Yarovoy,
E. Tajkhorshid,
B. Roux, and K. Schulten.
Calculation of the gating charge for the Kv1.2 voltage-activated potassium channel. *Biophysical Journal*, **98**:2189-2198, 2010.

350,000 atoms, 0.5 μ s


1M Atom Virus on TitanDev GPU





From 100,000 to 64,000,000 Atom MD 2013




100-million Atom Simulation 2014




**Simulated chromatophore with
101 protein complexes and
16,000 lipids**



**Tuning a
NAMD Run
on Blue
Waters**




Large is no problem. But ...



Molecular dynamics simulation of alpha-hemolysin with about 300,000 atoms; 10 million atom simulations are routine today, 200 million atom simulations are possible.



But long is still a problem! *biomolecular timescale and timestep limits*



(NSF center, Shaw Res.)

Preparing Your System for MD

Solvation

Biological activity is the result of interactions between molecules and occurs at the interfaces between molecules (protein-protein, protein-DNA, protein-solvent, DNA-solvent, etc).

*mitochondrial
bc1 complex*

Why model solvation?

- many biological processes occur in aqueous solution
- solvation effects play a crucial role in determining molecular conformation, electronic properties, binding energies, etc

How to model solvation?

- explicit treatment: solvent molecules are added to the molecular system
- implicit treatment: solvent is modeled as a continuum dielectric or so-called implicit force field



Preparing Your System for MD

Solvation

Biological activity is the result of interactions between molecules and occurs at the interfaces between molecules (protein-protein, protein-DNA, protein-solvent, DNA-solvent, etc).


*mitochondrial
bc1 complex*

Why model solvation?

- many biological processes occur in aqueous solution
- solvation effects play a crucial role in determining molecular conformation, electronic properties, binding energies, etc

How to model solvation?

- explicit treatment: solvent molecules are added to the molecular system
- implicit treatment: solvent is modeled as a continuum dielectric



Preparing Your System for MD

Solvation

Biological activity is the result of interactions between molecules and occurs at the interfaces between molecules (protein-protein, protein-DNA, protein-solvent, DNA-solvent, etc).


Why model solvation?

- many biological processes occur in aqueous solution
- solvation effects play a crucial role in determining molecular conformation, electronic properties, binding energies, etc

How to model solvation?

- explicit treatment: solvent molecules are added to the molecular system
- implicit treatment: solvent is modeled as a continuum dielectric


*mitochondrial
bc1 complex*



**(Usually periodic!
Avoids surface effects)**


From the Mountains to the Valleys

how to actually describe a protein



From the Mountains to the Valleys


a molecular dynamics tale




Cutting Corners

cutoffs, PME, rigid bonds, and multiple timesteps


- Nonbonded interactions require order N^2 computer time!
 - Truncating at R_{cutoff} reduces this to order $N R_{\text{cutoff}}^3$
 - Particle mesh Ewald (PME) method adds long range electrostatics at order $N \log N$, only minor cost compared to cutoff calculation.
- Can we extend the timestep, and do this work fewer times?
 - Bonds to hydrogen atoms, which require a 1fs timestep, can be held at their equilibrium lengths, allowing 2fs steps.
 - Long range electrostatics forces vary slowly, and may be evaluated less often, such as on every second or third step.
- Coarse Graining



Residue-Based Coarse-Grained Model



- Protein model uses two CG beads per residue
- One CG bead per side chain another for backbone





- Lipid model: MARTINI
- Level of coarse-graining: ~4 heavy atoms per CG bead
- Interactions parameterized based on experimental data and thermodynamic properties of small molecules

Peter L. Freddolino, Anton Arkhipov, Amy Y. Shih, Ying Yin, Zhongzhou Chen, and Klaus Schulten. **Application of residue-based and shape-based coarse graining to biomolecular simulations.** In Gregory A. Voth, editor, *Coarse-Graining of Condensed Phase and Biomolecular Systems*, chapter 20, pp. 299-315. Chapman and Hall/CRC Press, Taylor and Francis Group, 2008.

Nanodisc Assembly CG MD Simulation


- 10 μ s simulation
- Assembly proceeds in two steps:
 - Aggregation of proteins and lipids driven by the hydrophobic effect
 - Optimization of the protein structure driven by increasingly specific protein-protein interactions
- Formation of the generally accepted double-belt model for discoidal HDL



A. Shih, A. Arkhipov, P. Freddolino, and K. Schulten. *J. Phys. Chem. B*, 110:3674–3684, 2006; A. Shih, P. Freddolino, A. Arkhipov, and K. Schulten. *J. Struct. Biol.*, 157:579–592, 2007; A. Shih, A. Arkhipov, P. Freddolino, S. Sligar, and K. Schulten. *Journal of Physical Chemistry B*, 111: 11095–11104, 2007; A. Shih, P. Freddolino, S. Sligar, and K. Schulten. *Nano Letters*, 7:1692–1696, 2007.

Validation of Simulations

reverse coarse-graining and small-angle X-ray scattering




Reverse coarse-graining:


1. Map center of mass of the group of atoms represented by a single CG bead to that bead's location
2. MD minimization, simulated annealing with restraints, and equilibration to get all-atom structure

Small-angle X-ray scattering:

Calculated from reverse coarse-grained all-atom model and compared with experimental measurements



Shape-Based Coarse-Grained (CG) model





Peter L. Freddolino, Anton Arkhipov, Amy Y. Shih, Ying Yin, Zhongzhou Chen, and Klaus Schulten. **Application of residue-based and shape-based coarse graining to biomolecular simulations.** In Gregory A. Voth, editor, *Coarse-Graining of Condensed Phase and Biomolecular Systems*, chapter 20, pp. 299-315. Chapman and Hall/CRC Press, Taylor and Francis Group, 2008.

Virus Capsid Mechanics

Atomic Force Microscope


— Hepatitis B Virus —




Example: MD Simulations of the K⁺ Channel Protein

Ion channels are membrane-spanning proteins that form a pathway for the flux of inorganic ions across cell membranes.

Potassium channels are a particularly interesting class of ion channels, managing to distinguish with impressive fidelity between K⁺ and Na⁺ ions while maintaining a very high throughput of K⁺ ions when gated.




Setting up the system (1)



- retrieve the PDB (coordinates) file from the Protein Data Bank
- add hydrogen atoms using PSFGEN
- use psf and parameter files to set up the structure; needs to be better than available in Charmm to describe well the ions, e.g., K/Na ion selectivity or Ca^{++} ion
- minimize the protein structure using NAMD2


Setting up the system (2)



Simulate the protein in its natural environment: solvated lipid bilayer


Setting up the system (3)

Inserting the protein in the lipid bilayer



Automatic insertion into the lipid bilayer leads to big gaps between the protein and the membrane \Rightarrow long equilibration time required to fill the gaps. Solution: manually adjust the position of lipids around the protein. Employ constant (lateral and normal) pressure control.

The system



solvent

Kcsa channel protein (in blue) embedded in a (3:1) POPE/POPG lipid bilayer. Water molecules inside the channel are shown in vdW representation.

solvent

Simulating the system:


Free MD

Summary of simulations:


- protein/membrane system contains 38,112 atoms, including 5117 water molecules, 100 POPE and 34 POPG lipids, plus K^+ counterions
- CHARMM26 forcefield
- periodic boundary conditions, PME electrostatics
- 1 ns equilibration at 310K, NpT
- 2 ns dynamics, NpT

Program: NAMD2


Platform: Cray T3E (Pittsburgh Supercomputer Center) or local computer cluster; choose ~1000 atoms per processor.



Simulation of Ion Conduction (here for Kv1.2)



Voltage-gated Potassium Ion Channel



Voltage-gated Potassium Ion Channel

