



Molecular Dynamics of Proteins



Fibronectin III_1, a mechanical protein that glues cells together in wound healing and in preventing tumor metastasis



ATPase, a molecular motor that synthesizes the body's weight of ATP a day



AQP filtering a bath tub of the body's water a day




A ternary complex of DNA, lac repressor, and CAP controlling gene expression


Molecular Dynamics of Proteins



Fibronectin III_1, a mechanical protein that glues cells together in wound healing and in preventing tumor metastasis




ATPase, a molecular motor that synthesizes the body's weight of ATP a day



A ternary complex of DNA, lac repressor, and CAP controlling gene expression

The Molecular Dynamics Simulation Process



Classical Dynamics *at 300K*

Energy function: $U(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) = U(\vec{R})$

used to determine the force on each atom:

$$m_i \frac{d^2 \vec{r}_i}{dt^2} = \vec{F}_i = -\vec{\nabla} U(\vec{R})$$

yields a set of $3N$ coupled 2^{nd} -order differential equations that can be propagated forward (or backward) in time.

Initial coordinates obtained from crystal structure, velocities taken at random from Boltzmann distribution.

Maintain appropriate temperature by adjusting velocities.

Langevin Dynamics

feel the noise

Langevin dynamics deals with each atom separately, balancing a small friction term with Gaussian noise to control temperature:

$$m \ddot{\vec{r}} = \vec{F}(\vec{r}) - \gamma m \dot{\vec{r}} + \vec{R}(t)$$

$$\langle \vec{R}(t) \cdot \vec{R}(t') \rangle = 6k_B T \gamma \delta(t - t')$$

Classical Dynamics

discretization in time for computing

$$m_i \frac{d^2 \vec{r}_i}{dt^2} = \vec{F}_i = -\vec{\nabla} U(\vec{R})$$

Use positions and accelerations at time t and the positions from time $t-\delta t$ to calculate new positions at time $t+\delta t$.

$$\begin{aligned} \mathbf{r}(t + \delta t) &\approx \mathbf{r}(t) + \mathbf{v}(t)\delta t + \frac{1}{2}\mathbf{a}(t)\delta t^2 \\ \mathbf{r}(t - \delta t) &\approx \mathbf{r}(t) - \mathbf{v}(t)\delta t + \frac{1}{2}\mathbf{a}(t)\delta t^2 \end{aligned} \quad +$$

“Verlet algorithm”




$$-\vec{\nabla} U(\vec{R}) / m_i$$


$$\mathbf{r}(t + \delta t) \approx 2\mathbf{r}(t) - \mathbf{r}(t - \delta t) + \mathbf{a}(t)\delta t^2$$

Potential Energy Function of Biopolymer

- Simple, fixed algebraic form for every type of interaction.
- Variable parameters depend on types of atoms involved.



Large is no problem. But ...




Molecular dynamics simulation of alpha-hemolysin with about 300,000 atoms



NCSA machine room

But long is!

biomolecular timescale and timestep limits



PDB Files


gives one the structure and starting position

- Simulations start with a crystal structure from the Protein Data Bank, in the standard PDB file format.
- PDB files contain standard records for species, tissue, authorship, citations, sequence, secondary structure, etc.
- We only care about the atom records...
 - atom name (N, C, CA)
 - residue name (ALA, HIS)
 - residue id (integer)
 - coordinates (x, y, z)
 - occupancy (0.0 to 1.0)
 - temp. factor (a.k.a. beta)
 - segment id (6PTI)
- No hydrogen atoms!
(We must add them ourselves.)



PSF Files

- Every atom in the simulation is listed.
- Provides all static atom-specific values:
 - atom name (N, C, CA)
 - atom type (NH1, C, CT1)
 - residue name (ALA, HIS)
 - residue id (integer)
 - segment id (6PTI)
 - atomic mass (in atomic mass units)
 - partial charge (in electronic charge units)
- What is not in the PSF file?
 - coordinates (dynamic data, initially read from PDB file)
 - velocities (dynamic data, initially from Boltzmann distribution)
 - force field parameters (non-specific, used for many molecules)




PSF Files

molecular structure (bonds, angles, etc.)



Bonds: Every pair of covalently bonded atoms is listed.

Angles: Two bonds that share a common atom form an angle.
Every such set of three atoms in the molecule is listed.



Dihedrals: Two angles that share a common bond form a dihedral.
Every such set of four atoms in the molecule is listed.



Impropers: Any *planar* group of four atoms forms an improper.
Every such set of four atoms in the molecule is listed.



Preparing Your System for MD Solvation

Biological activity is the result of interactions between molecules and occurs at the interfaces between molecules (protein-protein, protein-DNA, protein-solvent, DNA-solvent, etc).

Why model solvation?

- many biological processes occur in aqueous solution
- solvation effects play a crucial role in determining molecular conformation, electronic properties, binding energies, etc

How to model solvation?

- explicit treatment: solvent molecules are added to the molecular system
- implicit treatment: solvent is modeled as a continuum dielectric or so-called implicit force field

*mitochondrial
bcl complex*



Preparing Your System for MD Solvation

Biological activity is the result of interactions between molecules and occurs at the interfaces between molecules (protein-protein, protein-DNA, protein-solvent, DNA-solvent, etc).


Why model solvation?

- many biological processes occur in aqueous solution
- solvation effects play a crucial role in determining molecular conformation, electronic properties, binding energies, etc

How to model solvation?

- explicit treatment: solvent molecules are added to the molecular system
- implicit treatment: solvent is modeled as a continuum dielectric

*mitochondrial
bcl complex*



Preparing Your System for MD Solvation

Biological activity is the result of interactions between molecules and occurs at the interfaces between molecules (protein-protein, protein-DNA, protein-solvent, DNA-solvent, etc).


Why model solvation?

- many biological processes occur in aqueous solution
- solvation effects play a crucial role in determining molecular conformation, electronic properties, binding energies, etc

How to model solvation?

- explicit treatment: solvent molecules are added to the molecular system
- implicit treatment: solvent is modeled as a continuum dielectric


*mitochondrial
bcl complex*



(Usually periodic!
Avoids surface effects)


From the Mountains to the Valleys

how to actually describe a protein



From the Mountains to the Valleys

a molecular dynamics tale




NAMD: The Program we will Use

Development cost 1990 - 2007: \$20 million



NAMD Registrants
 19,995 Registrants (3336 NIH)
 4,111 Repeat Users
 NAMD 2.6 released Aug 2006
 4181 NAMD 2.6 users (742 NIH)




NAMD
 programmer
 J. Phillips
 Ph.D. UIUC
 Physics

- ◆ CRAY T3E (2000)
- ◆ APOLLO (2000)
- ◆ XT3 (2006)
- ◆ XT3 (2006)
- ◆ XT3 (2006)
- ◆ XT3 (2006)
- ◆ XT3 (2006)

IAPP

NAMD scales by 10³



STMV

"We haven't found a hard limit on scaling up the number of processors."
 -- Philip Blood and Greg Voth,
 Univ Utah

Commenting on NAMD performance for the PSC XT3 Cray

Molecular Dynamics Ensembles

Constant energy, constant number of particles (NE)

Constant energy, constant volume (NVE)

Constant temperature, constant volume (NVT)


Constant temperature, constant pressure (NPT)

Choose the ensemble that best fits your system and start the simulations, but use NE to check on accuracy of the simulation.


Cutting Corners

cutoffs, PME, rigid bonds, and multiple timesteps


- Nonbonded interactions require order N^2 computer time!
 - Truncating at R_{cutoff} reduces this to order $N R_{\text{cutoff}}^3$
 - Particle mesh Ewald (PME) method adds long range electrostatics at order $N \log N$, only minor cost compared to cutoff calculation.
- Can we extend the timestep, and do this work fewer times?
 - Bonds to hydrogen atoms, which require a 1fs timestep, can be held at their equilibrium lengths, allowing 2fs steps.
 - Long range electrostatics forces vary slowly, and may be evaluated less often, such as on every second or third step.
- Coarse Graining



Residue-Based Coarse-Grained Model



- Protein model uses two CG beads per residue
- One CG bead per side chain another for backbone




- Lipid model: MARTINI
- Level of coarse-graining: ~4 heavy atoms per CG bead
- Interactions parameterized based on experimental data and thermodynamic properties of small molecules

Peter L. Freddolino, Anton Arkhipov, Amy Y. Shi, Ying Yin, Zhongzhou Chen, and Klaus Schulten. **Application of residue-based and shape-based coarse graining to biomolecular simulations.** In Gregory A. Voith, editor, *Coarse-Graining of Condensed Phase and Biomolecular Systems*, chapter 20, pp. 299-315. Chapman and Hall/CRC Press, Taylor and Francis Group, 2008.

Nanodisc Assembly CG MD Simulation

- 10 μs simulation
- Assembly proceeds in two steps:
 - Aggregation of proteins and lipids driven by the hydrophobic effect
 - Optimization of the protein structure driven by increasingly specific protein-protein interactions
- Formation of the generally accepted double-belt model for discoidal HDL



Validation of Simulations

reverse coarse-graining and small-angle X-ray scattering




Reverse coarse-graining:


1. Map center of mass of the group of atoms represented by a single CG bead to that beads location
2. MD minimization, simulated annealing with restraints, and equilibration to get all-atom structure

Small-angle X-ray scattering:

Calculated from reverse coarse-grained all-atom model and compared with experimental measurements




Shape-Based Coarse-Grained (CG) model



- Fully automatic
- Number of CG beads is chosen by a user (we used ~200 atoms per CG bead)


Peter L. Freddolino, Anton Arkhipov, Amy Y. Shih, Ying Yin, Zhongzhou Chen, and Klaus Schulten. **Application of residue-based and shape-based coarse graining to biomolecular simulations.** In Gregory A. Voth, editor. *Coarse-Graining of Condensed Phase and Biomolecular Systems*, chapter 20, pp. 299-315. Chapman and Hall/CRC Press, Taylor and Francis Group, 2008.

Reversible and irreversible indentations



Steps in a Typical MD Simulation


1. Prepare molecule
 - Read in pdb and psf file
 - Usually requires setting up the system, e.g., solvation
 - Many tools available in VMD
2. Minimization
 - Reconcile observed structure with force field used ($T = 0$)
3. Heating
 - Raise temperature of the system
4. Equilibration
 - Ensure system is stable
5. Dynamics
 - Simulate under desired conditions (NVE, NpT, etc)
 - Collect your data
6. Analysis
 - Evaluate observables (macroscopic level properties)
 - Or relate to single molecule experiments




Example: MD Simulations of the K⁺ Channel Protein

Ion channels are membrane-spanning proteins that form a pathway for the flux of inorganic ions across cell membranes.

Potassium channels are a particularly interesting class of ion channels, managing to distinguish with impressive fidelity between K⁺ and Na⁺ ions while maintaining a very high throughput of K⁺ ions when gated.




Setting up the system (1)



- retrieve the PDB (coordinates) file from the Protein Data Bank
- add hydrogen atoms using PSFGEN
- use psf and parameter files to set up the structure
- minimize the protein structure using NAMD2


Setting up the system (2)



Simulate the protein in its natural environment: solvated lipid bilayer


Setting up the system (3)

Inserting the protein in the lipid bilayer



Automatic insertion into the lipid bilayer leads to big gaps between the protein and the membrane \Rightarrow long equilibration time required to fill the gaps. Solution: manually adjust the position of lipids around the protein. Employ constant (lateral and normal) pressure control.

The system



Simulating the system: Free MD


Summary of simulations:

- protein/membrane system contains 38,112 atoms, including 5117 water molecules, 100 POPE and 34 POPG lipids, plus K^+ counterions
- CHARMM26 forcefield
- periodic boundary conditions, PME electrostatics
- 1 ns equilibration at 310K, NpT
- 2 ns dynamics, NpT


Program: NAMD2

Platform: Cray T3E (Pittsburgh Supercomputer Center) or local computer cluster; choose ~1000 atoms per processor.

MD Results




RMS deviations for the KcsA protein and its selectivity filter indicate that the protein is stable during the simulation with the selectivity filter the most stable part of the system.



Temperature factors for individual residues in the four monomers of the KcsA channel protein indicate that the most flexible parts of the protein are the N and C terminal ends, residues 52-60 and residues 84-90. Residues 74-80 in the selectivity filter have low temperature factors and are very stable during the simulation.

Simulation of Ion Conduction (here for Kv1.2)



Theoretical and Computational Biophysics Group Developers



- focus on systems biology
- theoretical biophysics
- develops renewable energy
- focus on quantum biology
- computational biophysics
- guides bionanotechnology